

Immunogenicity Studies of MenB Vaccines in Adults

ACIP

October 19, 2016

Dan Granoff, M.D., FPIDS
Center for Immunobiology and Vaccine Development
UCSF Benioff Children's Hospital Oakland
5700 Martin Luther King, Jr. Way
Oakland CA 94609 U.S.A.

Background

- Men A, C, Y and W capsular based vaccines, minimal antigenic variability within a capsular group
- Disease-causing MenB strains, protein antigens have large variability in amino acid sequence and expression, which can affect susceptibility to serum bactericidal activity
- For vaccine MenB licensure, efficacy was inferred based on data against a limited number of “reference strains”
- Gaps in knowledge
 - Extent of protection against more diverse disease-causing strains
 - Effect of vaccination schedule (2 doses vs 3), and duration of protection

Three Separate MenB Immunogenicity Studies in Adults

- Not designed to provide comparison data between vaccines
- Independent of industry
- No relevant potential conflicts of interest – other than inventor on patents related to meningococcal vaccines (assigned to UCSF Benioff Children's Hospital)

Conclusions

- One month post dose 2, both vaccines elicited protective bactericidal antibody (titers $\geq 1:4$) against most strains
- Some strains are relatively resistant to bactericidal activity despite prediction of susceptibility by sequence analysis and antigen expression
- After dose 2, titers can decline within 4 to 6 months, especially for strains with low antigen expression

Study 1, MenB-4C*

Immunogenicity in Adults

Study Sites	Oxford Vaccine Clinic, UK (N=15)
	UCSF Benioff Children's Hospital Oakland (N=5)
Median age, years	29
Healthcare or lab worker, No. (%)	12 (60)
Vaccine Schedule, Mos.	2 doses, 0,1 or 0, 2

*Bexsero, GSK

Giuntini et al,
Clinical Vaccine Immunol, 2016
In press

MenB-4C: Four Antigens




FHbp
(sub-family
B)



NHba




NadA




PorA
(P1.4)

≥ 4 -Fold Increases in Serum Bactericidal Antibody Titer to MenB-4C*




**Studies 1 and 2, data from
FDA package Insert;
Current study, N=20**

Sub-family B FHbp sequence variants (ID)



Adapted from, Granoff et al, J Infect Dis 2015


Serum Bactericidal Antibody Responses of Healthy Adults Immunized with 2 Doses of MenB-4C



Giuntini et al,
Clin Vac Immunol 2016, in press

MenB-4C has Sub-family B FHbp

Serum Bactericidal Antibody Responses of Adults Immunized with 2 Doses of MenB-4C*



**≥4-fold titer
Increases (1 mo
Post-dose 2/Pre)**

***, + or ++ NHba
†, + or ++ NadA**

**MenB-4C has Sub-
family B FHbp**

Study 2, MenB-FHbp*


Immunogenicity in Adults

Study Sites	UCSF Benioff Children's Hospital Oakland (N=12) University of Massachusetts Medical Center (N=5)
Median age, years	40
Healthcare or lab worker, No. (%)	17 (100)
Vaccine Schedule, Mos.	3 doses, 0, 2 and 6


*Trumenba, Pfizer

*Lujan et al, International
Pathogenic Neisseria
Conference (IPNC), 2016

MenB-FHbp (Trumenba, Pfizer)




FHbp Sub-family A and B




Serum Bactericidal Antibody Responses of Adults Immunized with MenB-FHbp: 0,2 Mo Schedule*

1 Mo Post Two Doses

FHbp Sub-Family A Strains



Pfizer Data




Data
from
Package
Insert

≥4-fold Increases in titers
(1 mo post-dose 2/Pre)


*Lujan et al, IPNC 2016

Serum Bactericidal Antibody Responses of Adults Immunized with Two Doses of MenB-FHbp: 0,2 Mos

≥ 4 -Fold titer Increases (1 Mo Post dose 2/Pre titers)



Pfizer Data





Subjects with <4-fold Increases in SBA Titer 1
Mo After Vaccination Can have “Protective”
Serum Bactericidal Titers ($\geq 1:4$)

Bactericidal Titers for a Representative
Strain (Quebec 2013, Hyperendemic)


FHbp sub-family B (ID 15)
Also expresses NHba

Serum Bactericidal Antibody Responses: Quebec 2013

Two Doses, Separated by 1 to 2 Mos.



Third Dose



PD2, post dose 2
PD3, post dose 3

Subjects with pre-titers $\geq 1:4$ excluded


Study 3, MenB-4C Immunogenicity in Santa Clara University Students

- January 31-February 3, 2016, 3 cases (2 culture confirmed), ST 32 clonal complex
- MenB-4C vaccination clinics
 - Feb 4-8, 4,921 persons received first vaccine dose
 - March 18 and April 6-8, 4,731 persons immunized (most receiving second dose)
- Sera collected May 23-25, from 246 students - 0 dose, N=52; 1 dose, N=91; and 2 doses, N=101
- Measured serum bactericidal against the college outbreak strains

Princeton University Outbreak Strain

- 9 cases, 1 death
- Outbreak strain, expressed two MenB-4C antigens, FHbp subfamily B and NHba
- Previous study
 - 66% of students immunized with 2 doses of MenB-4C, had protective serum bactericidal titers $\geq 1:4$ (Basta et al, NEJM 2016)

Prevalence of Serum Bactericidal Titers of Santa Clara Univ. Students Immunized with MenB-4C



Strain expresses two MenB-4C antigens, FHbp subfamily B (ID 276) and NHba

Interim results

MenB-4C Bactericidal Activity

- Complex vaccine, 4 antigens capable of eliciting serum bactericidal activity
- Antigen-specific reference strains are more susceptible to vaccine-induced bactericidal antibodies than many circulating disease-causing strains
- Some strains are relatively resistant to bactericidal activity despite prediction of susceptibility by sequence analysis and antigen expression
- Great majority of subjects have titers of $\geq 1:4$ at 1 mo post dose 2 against most strains
- Titers decline by 4 to 6 mos, especially against low FHbp-expressing strains

MenB-FHbp, 2 Injections, Separated by 2 Mos

- At 1 Mo post dose 2, similar ≥ 4 -fold increases in serum bactericidal activity as MenB-4C (against most strains)
- Great majority of subjects have titers of $\geq 1:4$ at 1 mo post dose 2 against most strains
- Titers decline by 4 mos, especially against low FHbp-expressing strains
- Recommended third dose of MenB-FHbp at 6 months boosts titers
- Additional data on antibody persistence needed (should include strains relatively resistant to vaccine-induced bactericidal activity)

Limitations of Current Studies

- Not designed to provide comparison data between vaccines
- Relatively few sera 4 to 6 months post-dose 2
- Serum bactericidal assays used research assays (not FDA validated)
- Data on sera from Santa Clara University study are antibody prevalence and interim analyses

MenB-4C
Immunogenicity Study

**UCSF Benioff Children's
Hospital, Oakland**

Eduardo Lujan, M.S.
Serena Giuntini, Ph.D.

Oxford Vaccine Group
University of Oxford, U.K

Malick M. Gibana, MRCP
Christina Dold, Ph.D.
Christine S. Rollier, Ph.D.
Andrew J. Pollard, FRCPCH PhD

MenB-FHbp
Immunogenicity Study

**UCSF Benioff Children's
Hospital, Oakland**

Eduardo Lujan, M.S.
Serena Giuntini, Ph.D.
Elizabeth Partridge, M.D.
Betty Flores R.N., PNP-BC, CCRP
Lisa H. Lavrisha, RN, PNP

**University of Massachusetts
Medical Center, Worcester**

Sanjay Ram, M.D.

Outbreak Strains, CDC

Santa Clara University

MenB-4C Serology Study

**UCSF Benioff Children's
Hospital, Oakland**

Flor Alicia Gowans, B.S.

Eduardo Lujan, M.S.

Santa Clara University
Jillandra C. Rovaris, Ph.D.

**California Department of Public
Health Immunization Branch**

Kathleen Winter, PhD, MPH

**Santa Clara County Public
Health Department**

George Han, MD, MPH

Wistar Institute, Philadelphia

Qin Liu, Ph.D.